



Anthelmintic activity of sesquiterpene lactones from forage chicory against *Ascaris suum*

Williams, Andrew R. ; Pena-Espinoza, Miguel Angel; Hansen, Tina V.A. ; Boas, Ulrik; Thamsborg, Stig M.

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Williams, A. R., Pena-Espinoza, M. A., Hansen, T. V. A., Boas, U., & Thamsborg, S. M. (2015). *Anthelmintic activity of sesquiterpene lactones from forage chicory against Ascaris suum*. Abstract from 6th Conference of the Scandinavian-Baltic Society for Parasitology, Uppsala, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ANTHELMINTIC ACTIVITY OF SESQUITERPENE LACTONES FROM FORAGE CHICORY AGAINST *ASCARIS SUUM*

Andrew R. Williams¹, Miguel Peña-Espinoza², Tina V.A. Hansen¹, Ulrik Boas², Stig M. Thamsborg¹

¹*Department of Veterinary Disease Biology, University of Copenhagen, Denmark.*

²*National Veterinary Institute, Technical University of Denmark, Denmark.*

arw@sund.ku.dk

Introduction

New control options for parasitic nematodes are needed due to the threat of drug resistance and consumer demand for organic animal products. Chicory is a nutritious forage that can be used for grazing outdoor-reared pigs and ruminants and may have anthelmintic effects, potentially due to high concentrations of sesquiterpene lactones (SL). Here, we investigated the effects of SL extracts from chicory on *in vitro* survival and glutathione-S-transferase (GST) activity of the highly prevalent pig parasite *Ascaris suum*.

Materials and Methods

Chicory leaves from two cultivars (Spadona and Puna II) were extracted with methanol and SL purified by solid-phase extraction. UPLC-MS was used for analysis of the extracts. *A. suum* third-stage larvae (L3) were produced by *in vitro* hatching of eggs, whilst L4 were isolated from the gut of infected pigs. Native *A. suum* GST protein was purified and activity was assessed by the CNDB assay.

Results

Incubation in the SL extracts dramatically reduced the *in vitro* survival of *A. suum* L3 and L4. However, the effect was cultivar-dependent, with Spadona extracts more potent than Puna II, consistent with distinct SL profiles observed between cultivars. Spadona-SL also strongly inhibited the activity of *A. suum* GST, suggesting that the anthelmintic mechanism may involve accumulation of toxic reactive oxygen.

Conclusion

If the observed anthelmintic effects are confirmed *in vivo*, chicory has potential to be used as an alternative control for *A. suum* in outdoor pigs, although selection of appropriate cultivars will be important. On-going experiments are investigating the mechanism(s) of anthelmintic action.